

1. The number of solutions of the system of equations

$$2x + y - z = 7; x - 3y + 2z = 1; x + 4y - 3z = 5$$

(A) 0

(B) 2

(C) 3

(D) 1

2. For non-zero numbers p, q, r, a, b, c , if $\begin{vmatrix} pa & qb & rc \\ qc & ra & pb \\ rb & pc & qa \end{vmatrix} = pqr \begin{vmatrix} a & b & c \\ c & a & b \\ b & c & a \end{vmatrix}$ then

(A) $pqr = 1$

(B) $p + q + r = 1$

(C) $p + q + r = 0$

(D) $pqr = 0$

3. Let $\begin{vmatrix} x & 2 & x \\ x^2 & x & 6 \\ x & x & 6 \end{vmatrix} = Ax^4 + Bx^3 + Cx^2 + Dx + E$, then the value of $9A - 4B + 3C + 5D + 6E$ is

(A) 36

(B) 38

(C) 35

(D) 37

Space for Rough Work

4. If $A = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$, then the matrix A^{2014} is same as

- (A) A
- (B) $-A$
- (C) I
- (D) $-I$

5. If $A = \begin{vmatrix} -1 & 2 & 0 \\ 3 & 1 & 5 \\ -1 & 2 & -1 \end{vmatrix}$, then $|\text{adj}(\text{adj}A)|$ is

- (A) 1492
- (B) 1592
- (C) 1694
- (D) 2401

6. The value of $\sec^2(\tan^{-1}2) + \text{cosec}^2(\cot^{-1}3)$ is

- (A) 5
- (B) 20
- (C) 10
- (D) 15

Space for Rough Work

7. Given that $\sin \alpha + \sin \beta = p$ and $\cos \alpha - \cos \beta = q$, then the value of $\cos(\alpha - \beta)$ is

(A) $\frac{p^2 - q^2}{p^2 + q^2}$

(B) $\frac{2pq}{p^2 + q^2}$

(C) $\frac{2pq}{p^2 - q^2}$

(D) $\frac{p^2 + q^2}{p^2 - q^2}$

8. The value of $\sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ$ is

(A) $\frac{1}{2}$

(B) $\frac{1}{4}$

(C) 2

(D) 4

9. If $x = \sin^2\theta + \cos^4\theta$, then for all values of θ , the interval x belongs to is

(A) $0 \leq x \leq 1$

(B) $1 \leq x \leq 2$

(C) $\frac{3}{4} \leq x \leq 1$

(D) $\frac{1}{4} \leq x \leq \frac{1}{2}$

Space for Rough Work

10. The equality $\cot^{-1}\alpha = \tan^{-1}\frac{1}{\alpha}$ holds good only when
- (A) $\alpha = 0$
 - (B) $|\alpha| \leq 1$
 - (C) $\alpha < 0$
 - (D) $\alpha > 0$
11. The approximate value of $\tan^{-1}(1.001)$ is
- (A) $\frac{\pi}{4} + 0.1$
 - (B) $\frac{\pi}{4} + 0.005$
 - (C) $\frac{\pi}{4} + 0.002$
 - (D) $\frac{\pi}{4} + 0.0005$
12. The roots of $x^2 - 2\sqrt{3}x + 2 = 0$ represent the lengths of two sides of a triangle and if the angle between these sides is 60° , then the perimeter of the triangle is
- (A) $3 + 2\sqrt{6}$
 - (B) $2\sqrt{3} + 2\sqrt{6}$
 - (C) $2\sqrt{3} + \sqrt{6}$
 - (D) $3\sqrt{2} + 6$

Space for Rough Work

13. The area bounded by the ellipse $\frac{x^2}{175} + \frac{y^2}{343} = 1$ is
- (A) $150\sqrt{7}\pi$
 - (B) 115π
 - (C) 200π
 - (D) 245π
14. If $x - 2y - a = 0$ is a chord of the parabola $y^2 = 4ax$, then its length is given by
- (A) $10a$
 - (B) $20a$
 - (C) $30a$
 - (D) $40a$
15. A pair of straight lines is given by $x^2(\sin^2\alpha - 1) + y^2\cos^2\alpha - yx \cos^2\alpha = 0$, the angle between them is given by
- (A) π
 - (B) $\frac{\pi}{4}$
 - (C) $\frac{\pi}{2}$
 - (D) $\frac{2\pi}{3}$

Space for Rough Work

16. If the projection of a line segment on x , y and z axes are 4, 2, $\sqrt{21}$ respectively, then length of the line segment is
- (A) $6 + \sqrt{21}$
- (B) $\sqrt{41}$
- (C) $4 + 2\sqrt{21}$
- (D) $\sqrt{43}$
17. The equations of common tangents to the circle $x^2 + y^2 = 8$ and parabola $y^2 = 16x$ are
- (A) $x = \pm (y + 2)$
- (B) $x = \pm (y + 4)$
- (C) $y = \pm (x + 2)$
- (D) $y = \pm (x + 4)$
18. If p , q , r are in Arithmetic Progression, then $px + qy + r = 0$ represents a
- (A) point
- (B) single line
- (C) family of concurrent lines
- (D) family of circles

Space for Rough Work

19. The tangents drawn at any point on these two curves $3x^2y - y^3 - 2 = 0$ and $x^3 - 3xy^2 + 2 = 0$ cut at
- (A) 90°
- (B) 60°
- (C) 45°
- (D) 30°
20. The smaller of the two areas enclosed between the ellipse $\frac{x^2}{4} + \frac{y^2}{16} = 1$ and the line $\frac{x}{2} + \frac{y}{4} = 1$ is
- (A) $2\pi - 4$
- (B) $2\left(\pi - \frac{1}{2}\right)$
- (C) $2\pi^2$
- (D) $\frac{\pi^2}{8}$
21. For some natural number n , if $\sum n = 55$, then $\sum n^2$ is
- (A) 3125
- (B) 605
- (C) 1025
- (D) 385

Space for Rough Work

22. If the sum of two of the roots of $x^3 + ax^2 + bx + c = 0$ is zero, then the value of ab is
- (A) $2c$
(B) $3c$
(C) $-c$
(D) c
23. The value of the sum ${}^{18}C_2 + {}^{18}C_4 + {}^{18}C_6 + \dots + {}^{18}C_{18}$ is
- (A) $2^{17} - 1$
(B) $2^{18} - 1$
(C) $2^{19} - 1$
(D) 2^{18}
24. The sum of the series $1 + \frac{x}{2} + \frac{x(x-1)}{2.4} + \frac{x(x-1)(x-2)}{2.4.6} + \dots$ to ∞ is
- (A) $\left(\frac{3}{4}\right)^x$
(B) $\left(\frac{4}{3}\right)^x$
(C) $\left(\frac{3}{2}\right)^x$
(D) $\left(-\frac{3}{2}\right)^x$

Space for Rough Work

25. The value of $5^{\frac{1}{2}} \cdot 5^{\frac{1}{4}} \cdot 5^{\frac{1}{8}} \dots$ to ∞ is

- (A) 25
- (B) 5
- (C) 125
- (D) $\frac{1}{5}$

26. The product of all n^{th} roots of unity ($n > 1$) is

- (A) 1
- (B) $(n)^{n-1}$
- (C) 0
- (D) $(-1)^{n-1}$

27. If α, β, γ are the direction cosines of a line then for some real number c , the value of $c[\cos 2\alpha + \cos 2\beta + \cos 2\gamma]$ is

- (A) $-c$
- (B) 0
- (C) $2c$
- (D) c

Space for Rough Work

28. If $\vec{a} = 3\hat{i} - 5\hat{j}$; $\vec{b} = 6\hat{i} + 3\hat{j}$ and $\vec{c} = \vec{a} \times \vec{b}$, then $|\vec{a}| : |\vec{c}| : |\vec{b}| =$

(A) $34 : 39 : 45$

(B) $39 : 35 : 34$

(C) $\sqrt{34} : 39 : \sqrt{45}$

(D) $\sqrt{34} : \sqrt{39} : \sqrt{45}$

29. If \vec{a}, \vec{b} are non-collinear vectors and x, y are scalars such that

$$(2\vec{a} - \vec{b})x + (2\vec{b} - \vec{a})y + (\vec{a} + 2\vec{b}) = \vec{0}, \text{ then}$$

(A) $x = -\frac{4}{3}, y = -\frac{5}{3}$

(B) $x = -\frac{4}{3}, y = \frac{5}{3}$

(C) $x = 0, y = 4$

(D) $x = \frac{5}{3}, y = \frac{4}{3}$

30. If ABCD is a square, then $\vec{AB} + 2\vec{BC} + 3\vec{CD} + 4\vec{DA}$ is

(A) $5\vec{CA}$

(B) $2\vec{CA}$

(C) $3\vec{CA}$

(D) $8\vec{CA}$

Space for Rough Work

31. The direction cosines of two lines that are at right angles are l_1, m_1, n_1 and l_2, m_2, n_2 , then the direction cosines of a line which is perpendicular to both these lines are
- (A) $l_1 + kl_2, m_1 - km_2, n_1 + kn_2$
- (B) $m_1n_2 - m_2n_1, n_1l_2 - n_2l_1, l_1m_2 - l_2m_1$
- (C) $l_1 - l_2, m_1 - m_2, n_1 - n_2$
- (D) $l_1 + l_2, m_1 + m_2, n_1 + n_2$
32. If $|\vec{a}| = 4$; $|\vec{b}| = 2$ and the angle between \vec{a} and \vec{b} is $\frac{\pi}{6}$, then $|\vec{a} \times \vec{b}|$ is
- (A) 3
- (B) 4
- (C) 16
- (D) 9
33. If the vectors $\vec{a} = 3\vec{i} + 6\vec{j} + 2\vec{k}$ and \vec{b} are collinear and $|\vec{b}| = 28$, then $\vec{b} =$
- (A) $\pm 3 (2\vec{i} + 6\vec{j} + \vec{k})$
- (B) $\pm 4 (3\vec{i} + 6\vec{j} + 2\vec{k})$
- (C) $\pm 28 (3\vec{i} + 6\vec{j} + 2\vec{k})$
- (D) $\pm 2 (3\vec{i} + 6\vec{j} + 2\vec{k})$

Space for Rough Work

34. For three vectors, $|\vec{a}| = |\vec{b}| = |\vec{c}|$, angle between each pair of these vectors is $\frac{\pi}{3}$ and

$$|\vec{a} + \vec{b} + \vec{c}| = \sqrt{6}, \text{ then } |\vec{a}| \text{ is}$$

- (A) 1
- (B) $\sqrt{3}$
- (C) -1
- (D) 2

35. The value of the limit $\lim_{x \rightarrow 0} \frac{\log(1+ax) - \log(1-bx)}{x}$ is

- (A) $a - b$
- (B) $a + b$
- (C) ab
- (D) $\frac{a}{b}$

36. If the function $x = x(y)$ is defined as $x = e^{y+e^y+e^{y+\dots\infty}}$ then $\frac{dy}{dx}$ is given by

- (A) $1 + x$
- (B) $\frac{1}{x}$
- (C) $\frac{1}{x} - 1$
- (D) $\frac{x}{1+x}$

Space for Rough Work

37. The value of the integral $\int e^{2x} \left(\frac{1 + \sin 2x}{1 + \cos 2x} \right) dx$ is

(A) $\frac{1}{2} e^{2x} \tan 2x + c$

(B) $\frac{1}{2} e^{2x} \tan x + c$

(C) $\frac{1}{2} e^{2x} \sin 2x + c$

(D) $\frac{1}{2} e^{2x} \cos 2x + c$

38. If $\int_m^{m+1} f(x) dx = m^2$, where $m \in \mathbb{Z}$, then $\int_{-1}^3 f(x) dx$ is

(A) 54

(B) 16

(C) 6

(D) 36

Space for Rough Work

39. If $f(\theta) = \left| \begin{array}{cc} \sec \theta & \cos \theta \\ \cos^2 \theta & \cos^2 \theta \end{array} \right|$, then the value of the definite integral $\int_0^{\pi/2} f(\theta) d\theta$ is

(A) $\frac{1}{5}$

(B) $\frac{1}{4}$

(C) $\frac{1}{3}$

(D) $\frac{1}{2}$

40. The solution of $\frac{dy}{dx} = \frac{1}{y^2 + \sin y}$, $y \neq 0$, with an arbitrary constant c is

(A) $x = y^3 - \cos^2 y + c$

(B) $x = y - \cos y + c$

(C) $x = \frac{y^3}{3} - \cos y + c$

(D) $x = y^2 - \frac{\cos y}{3} + c$

Space for Rough Work

41. The differential equation governing the solution $ax^2 - by^2 = 16$ is $y \left(\frac{d^2y}{dx^2} \right) + \left(\frac{dy}{dx} \right)^2 =$

(A) $y \frac{dy}{dx}$

(B) $\left(\frac{1}{x} \right) \frac{dy}{dx}$

(C) $\left(\frac{y}{x} \right) \frac{dy}{dx}$

(D) $\left(\frac{x}{y} \right) \frac{dy}{dx}$

42. If $\frac{dy}{dx} = u^2$, where $u = 4x + y + 1$ then,

(A) $2 \tan^{-1} \left(\frac{u}{2} \right) = x + c$

(B) $\tan^{-1} \left(\frac{u}{2} \right) = 2(x + c)$

(C) $y = 2 \tan^{-1} x + c$

(D) $\frac{u^2}{2} = \tan^{-1} xy + c$

Space for Rough Work

43. The order and degree of the differential equation $\sqrt[3]{\frac{dy}{dx}} \sqrt{\frac{d^3y}{dx^3}} = 4$ is

(A) 2, 3

(B) 3, 6

(C) 3, 2

(D) 3, 1

44. If $y = (1234)e^{11x} + (5678)e^{-11x}$ then $\frac{d^2y}{dx^2}$ is equal to

(A) 1234y

(B) 5678y

(C) 121y

(D) 1331y

45. The differential equation that represents the family of lines $ax + by + c = 0$ is

(A) $\frac{dy}{dx} = 0$

(B) $x + y \frac{dy}{dx} + \frac{d^2y}{dx^2} = 0$

(C) $\frac{d^2y}{dx^2} = 0$

(D) $y = x \frac{dy}{dx} + c$

Space for Rough Work

46. If $P(A) = x$, $P(B) = 2x$, $P(A \cap B) = \frac{1}{2}$, $P(\bar{A} \cap \bar{B}) = \frac{2}{3}$, then the value of x is
- (A) $\frac{5}{18}$
- (B) $\frac{5}{36}$
- (C) $\frac{6}{36}$
- (D) $\frac{11}{36}$
47. A die is rolled 3 times, the probability of getting a number larger than the previous number each time is
- (A) $\frac{5}{54}$
- (B) $\frac{1}{18}$
- (C) $\frac{13}{216}$
- (D) $\frac{23}{216}$
48. If Ramu and Raju can solve 80% and 60% respectively of the problems in a book, what is the probability that at least one of them will solve “the problem selected at random” from the book.
- (A) 0.92
- (B) 0.86
- (C) 0.68
- (D) 0.94

Space for Rough Work

49. The dual of the statement $p \vee (q \vee r) \equiv (p \vee q) \vee r$ is
- (A) $p \wedge (q \vee r) \equiv (p \wedge q) \vee r$
 - (B) $p \wedge (q \wedge r) \equiv (p \wedge q) \wedge r$
 - (C) $p \vee (q \wedge r) \equiv (p \wedge q) \vee r$
 - (D) None of these
50. A function is defined as $f(x) = \frac{kx}{x+1}$, $x \neq -1$, then for what value of k is $f(f(x)) = x$
- (A) -1
 - (B) 1
 - (C) $\sqrt{3}$
 - (D) $-\sqrt{2}$
51. The maximum value of $1 + 8 \sin^2\theta^2 \cos^2\theta^2$ is
- (A) 0
 - (B) -8
 - (C) 3
 - (D) 10

Space for Rough Work

52. The coefficient of x^4 in the expansion of $\frac{2x + 1 + 3x^2}{e^x}$ is
- (A) $-\frac{36}{15}$
- (B) $\frac{29}{24}$
- (C) $\frac{3}{2}$
- (D) $-\frac{8}{5}$
53. A set S has 5 distinct elements. Then the number of distinct one-one functions that can be defined from S to S is
- (A) 32
- (B) 2^{25}
- (C) 120
- (D) 5^5
54. The digit in the units place of $1! + 2! + 3! + 4! + \dots + n!$, where $n > 4$ is
- (A) 1
- (B) 2
- (C) 3
- (D) 4

Space for Rough Work

55. For $n \in \mathbb{N}$, $6^n - 5n - 1$ is always divisible by

- (A) 50
- (B) 25
- (C) 75
- (D) 125

56. The quadratic equation whose roots are p and q where

$$p = \lim_{x \rightarrow 0} \frac{3 \sin x - 4 \sin^3 x}{x} \quad \text{and} \quad q = \lim_{x \rightarrow 0} \frac{2 \tan x}{1 - \tan^2 x} \text{ is}$$

- (A) $x^2 + 5x + 6 = 0$
- (B) $x^2 + 3x + 2 = 0$
- (C) $x^2 - 5x + 6 = 0$
- (D) $x^2 - 3x + 2 = 0$

57. If $(1 + i)^{100} = 2^{49} (x + iy)$, then $x^2 + y^2$ is equal to

- (A) 0
- (B) 32
- (C) 16
- (D) 4

Space for Rough Work

58. For the complex number $i, i^{4n} + i^{4n+1} + i^{4n+2} + i^{4n+3} + i^{4n+4} + i^{4n+6}$ is
- (A) 16
 (B) 4
 (C) 1
 (D) 0
59. In the expansion of $\frac{(2-x)(2+x)}{(1-x)(1+x)}, |x| < 1$, the term that is independent of x is
- (A) 3
 (B) 4
 (C) 5
 (D) 2
60. For three real numbers a, b, c with $a \neq 6$; if $\begin{vmatrix} a & 2b & 2c \\ 3 & b & c \\ 4 & a & b \end{vmatrix} = 0$, then $abc =$
- (A) $a + b + c$
 (B) $ab + b - c$
 (C) 0
 (D) b^3

Space for Rough Work

Space For Rough Work